

GLAUCOMA



What is glaucoma?

Glaucoma is one of the leading causes of blindness in the United States, especially for older people. But loss of sight from glaucoma is often preventable if you get treatment early enough. Glaucoma is a disease of the optic nerve. The optic nerve carries the images we see to the brain. It is made up of a huge number of nerve fibers, like an electric cable containing a huge number of wires. Glaucoma can damage nerve fibers, causing blind spots to develop. Usually people don't notice these blind areas until much optic nerve damage has already occurred. If the entire nerve is destroyed, blindness results. Early detection and treatment by your eye care specialist are the keys to preventing optic nerve damage and blindness from glaucoma.

What causes glaucoma?

The complete mechanism of Glaucoma is not fully understood. What follows is a condensed version of what is known, but other factors likely play a role too.

Clear liquid, called the aqueous humor, circulates inside the eye. A small amount of this fluid is produced constantly, and an equal amount flows out of the eye through a microscopic drainage system. (This liquid is not part of the tears on the outer surface of the eye.) You can think of the flow of aqueous fluid as a sink with the faucet turned on all the time. If the "drainpipe" gets clogged, water collects in the sink and the sink may overflow. Because the eye is a closed structure, the excess fluid cannot overflow if the drain is clogged. If the drainage

area of the eye is blocked, the fluid pressure within the inner eye may increase, which can damage the optic nerve.

What are the different types of glaucoma?

Chronic open-angle glaucoma

This is the most common form of glaucoma in the United States. It occurs as a result of aging. The drainage angle of the eye becomes less efficient with time, and pressure within the eye gradually increases.

If this increased pressure results in optic nerve damage, it is known as chronic open-angle glaucoma. Over 90% of adult glaucoma patients have this type of glaucoma.

Chronic open-angle glaucoma damages vision so gradually and painlessly that you are not aware of trouble until the optic nerve is already badly damaged.

Angle-closure glaucoma

Sometimes the drainage angle of the eye may become completely blocked. It is as though a sheet of paper floating near a drain suddenly drops over the opening and blocks the flow out of the sink. In the eye, the iris (the part that makes eyes blue or brown) may act like the sheet of paper closing off the drainage angle. When eye pressure builds up suddenly, it is called acute angle-closure glaucoma.



Our doctors (from left): Matt Oliva, MD • Paul Imperia, MD • Paul Jorizzo, MD
Kimberly Hoyt, OD • Pete Spittelie, MD • Michael Hoyt, OD

You can depend on us.

We're a full-service eye care practice assisted by an exceptionally qualified staff. Medical Eye Center was founded in 1921 on the principle of preserving the health of our patients and the quality of their vision through comprehensive eye care and surgical services.



MEDICAL EYE CENTER

Paul Jorizzo, MD • Paul Imperia, MD
Pete Spittelie, MD • Matt Oliva, MD
Kimberly Hoyt, OD • Michael Hoyt, OD

Medford Grants Pass

1333 E. Barnett Road
Medford, OR 97504

Phone: 541-779-4711
Fax: 541-779-0796

881 N.E. 7th Street
Grants Pass, OR 97526

Phone: 541-476-6302
Fax: 541-476-1440

1-800-824-2688

www.medicaleyecenter.com



Symptoms of angle closure may include:

- Blurred vision
- Severe eye pain
- Headache
- Rainbow haloes around lights
- Nausea and vomiting

If you have any of these symptoms, call your eye care specialist immediately. Unless your doctor treats acute angle-closure glaucoma quickly, blindness can result. In some patients glaucoma has features of both the chronic open angle type and the acute angle closure type. This may be called chronic angle closure glaucoma or mixed mechanism glaucoma.

How is glaucoma detected?

Regular eye examinations by your eye care specialist are the best way to detect and treat glaucoma.

During a complete and painless examination, your eye care specialist will:

- Measure your intraocular pressure (tonometry)
- Evaluate any optic nerve damage (ophthalmoscopy)

Your Doctor may do additional testing including:

- Inspect the drainage angle of your eye (gonioscopy)
- Test the visual field of each eye (perimetry)

Some of these tests may not be necessary for every person. If you have glaucoma, you will need to repeat these tests on a regular basis, to determine if glaucoma is increasing over time.

Who is at risk for glaucoma?

High pressure alone does not mean that you have glaucoma. Your eye care specialist puts together many kinds of information to determine your risk for developing the disease.

The most important risk factors include:

- Age
- Family history of glaucoma
- Past injuries to the eyes
- African ancestry

Your eye care specialist will weigh all of these factors before deciding whether you need treatment for glaucoma, or whether you should be monitored closely as a glaucoma suspect. Glaucoma suspect means your risk of developing glaucoma is higher than normal, and you need to have regular examinations to detect early signs of damage to the optic nerve.

How is glaucoma treated?

Damage caused by glaucoma cannot be reversed. Eye drops, laser procedures and surgical operations are used to prevent or slow further damage from occurring. With any type of glaucoma, periodic examinations are very important to prevent vision loss. Because glaucoma can worsen without your being aware of it, your treatment may need to be changed over time.

Medicines

Glaucoma is often controlled with eye drops. These medications decrease eye pressure outflow, by slowing the production of aqueous fluids within the eye. For these medications to work, you must take them regularly and continuously. It is also important to tell all of your doctors about the eye medications you are using. Glaucoma medications can have side effects. You should notify your eye care specialist immediately if you think you may be experiencing side effects.

Some eye drops may cause:

- A stinging sensation
- Red eyes
- Changes in pulse and heartbeat
- Changes in energy level
- Changes in breathing (especially with asthma or emphysema)
- Headaches
- Blurred vision

Laser procedures

Laser surgery treatments may be effective for different types of glaucoma. The laser is usually used in one of two ways.

In open-angle glaucoma, the drain itself is treated. The laser is used to modify the drain (trabeculoplasty) to help control eye pressure. In angle-closure glaucoma, the laser creates a hole in the iris (iridotomy) to improve the flow of aqueous fluid to the drain.

Operative surgery

When operative surgery is needed to control glaucoma, your eye care specialist uses miniature instruments to create a new drainage channel for the aqueous fluid to leave the eye (trabeculectomy). The new channel helps to lower the pressure. Though serious complications of modern glaucoma surgery are rare, they can occur, as with any surgery. Surgery is recommended if your doctor feels that it is safer to operate than to allow optic nerve damage to continue.

What is your part in treatment?

Treatment for glaucoma requires teamwork between you and your doctor. Your eye care specialist can prescribe treatment for glaucoma, but only you can make sure you take your eye drops. Never stop taking or change your medications without first consulting your eye care specialist. Frequent eye examinations and tests are critical to monitor your eyes for any changes. Remember, it is your vision, and you must do your part to maintain it.

Loss of vision can be prevented

Regular yearly medical eye exams may help prevent unnecessary vision loss.